



Environment
Canada

Environnement
Canada

Canadian
Forestry
Service


Service
canadien des
forêts

Metric Single-tree Total Volume Tables for the Yukon Territory

M.R.C. Massie, G.H. Manning and K.R. McCloskey



Information Report BC-X-242
Pacific Forest Research Centre



DBH (cm)	Height (m)				
	4	6	8	10	12
4	.0024	.0037	.0050		
6	.0050	.0077	.0105	.0133	
8	.0084	.0130	.0177	.0225	.0273
10	.0126	.0195	.0266	.0337	.0410
12	.0176	.0272	.0370	.0470	.0572
14		.0360	.0490	.0623	.0758
16		.0459	.0626	.0795	.0967
18		.0569	.0775	.0985	.1198
20		.0690	.0940	.1194	.1452
22			.1118	.1420	.1727
24			.1310	.1664	.2024
26				.1926	.2342
28					.2681
30					
32					

**METRIC SINGLE-TREE TOTAL VOLUME TABLES
FOR THE YUKON TERRITORY**

M.R.C. Massie*, G.H. Manning and K.R. McCloskey***

* Nawitka Renewable Resource Consultants
Victoria, B.C.

** Pacific Forest Research Centre
Canadian Forestry Service
506 West Burnside Road
Victoria, B.C.
V8Z 1M5

Environment Canada
Canadian Forestry Service
Pacific Forest Research Centre

BC-X-242

1983

ABSTRACT

Metric single-tree volume tables are presented for the four major tree species in the Yukon Territory. Total inside-bark stem volume is shown as a function of total height and outside-bark diameter breast height.

RESUME

Des tables de volume en métrique pour un seul arbre sont présentées pour les quatre espèces importantes du territoire du Yukon. Le volume totale des tiges (sans écorces) est fonction de la hauteur totale de l'arbre et du diamètre a hauteur d'homme incluant l'écorce.

Environment Canada
Canadian Forestry Service
Pacific Forest Research Centre
506 West Burnside Road
Victoria, B.C.
V8Z 1M5

© Minister of Supply and Services Canada, 1983

ISSN 0705-3274
ISBN 0-662-12658-0
Cat. No. Fo46-17/242E

TABLE OF CONTENTS

Introduction	7
Methods	7
Results	7
References	7
Appendix A	15
Appendix B	16
Appendix C	17

LIST OF TABLES

Table		Page
1.	Cubic metre volume table for white spruce (all ages) Yukon Territory— Ecoregions 1, 2, 9, 11, 12.	3
2.	Cubic metre volume table for white spruce (all ages) Yukon Territory— Ecoregions 4, 5, 8.	4
3.	Cubic metre volume table for black spruce (all ages) Yukon Territory— Ecoregions 2, 5, 11, 12.	5
4.	Cubic metre volume table for black spruce (all ages) Yukon Territory— Ecoregions 4, 8, 9.	6
5.	Cubic metre volume table for lodgepole pine (all ages) Yukon Territory— Ecoregions 2, 5.	7
6.	Cubic metre volume table for lodgepole pine (all ages) Yukon Territory— Ecoregions 4, 12.	8
7.	Cubic metre volume table for trembling aspen (all ages) Yukon Territory— Ecoregions 2, 4, 5, 9, 11, 12.	9

INTRODUCTION

A study of tree component weights conducted by Nawitka Renewable Resource Consultants and funded by the Canadian Forestry Service destructively sampled 362 white spruce (*Picea glauca* (Moench) Voss), 300 black spruce (*Picea mariana* (Mill.) B.S.P.), 148 lodgepole pine (*Pinus contorta* Dougl.) and 175 trembling aspen (*Populus tremuloides* Michx.) in the Yukon between 1980 and 1982. These samples were randomly distributed across the accessible areas of ecoregions 1, 2, 4, 5, 8, 9, 11 and 12 (Oswald and Senyk 1977) on the basis of species representation (Gairns 1968). The exception was ecoregion 1, where only white spruce was sampled. Sample distribution is shown in Appendix A; ecoregion location is shown in Appendix B.

Sectioning of the sample trees during the biomass measurement allowed the calculation of whole tree volumes by Smalian's formula. Utilization of the accompanying field measurements permitted the development of equations to predict the volume for white spruce, lodgepole pine, black spruce and trembling aspen in each ecoregion.

METHODS

For each species the following procedure was used:

1. Height-diameter regressions were fitted to the sample trees in each ecoregion. Sample trees in each ecoregion were examined for seriously atypical trees by examination of scatter diagrams that related height to diameter. Three trees were rejected as being seriously atypical. Inspection of the height-diameter curves determined an initial grouping of ecoregions. Using these groupings, volume equations of the

type $V = aDBH^bHt^c$ were fitted using least squares regression. The groupings were subjected to covariance analysis to determine whether they represented an improvement over a single equation for the Yukon for each species. In all cases the result was significant at the 99% level. Statistics may be found in Appendix C.

2. Metric volume tables were constructed for each grouping by means of the derived logarithmic equation.
3. The statistical hypothesis of a single volume equation for mixed and pure stands was tested and shown to be valid.

RESULTS

The metric volume tables that follow show the total volume of the stem, inside bark, from germination to terminal bud. The diameter shown is standing DBH measured at 1.3 metres. Height is total height. There is no allowance for defect.

References

- Gairns, C.H. 1968. The Yukon Economy—Its Potential for Growth and Continuity: Volume VIII—References Study on Forest Resources. D. Wm. Carr & Associates for Dept. of Indian Affairs and Northern Development and the Government of Yukon Territory, Ottawa.
- Oswald, E.T. and J.P. Senyk. 1977. Ecoregions of Yukon Territory. Can. For. Serv., Pac. For. Res. Cent., Inf. Rep. BC-X-164.

Table 1. Cubic metre volume table for white spruce (all ages), Yukon Territory, Ecoregions 1, 2, 9, 11, and 12.

DBH (cm)	Height (m)																		Basis (No. of Trees)
	4	6	8	10	12	14	16	18	20	22	24	26	28	30	32	34	36	38	
4	.0024	.0037	.0051	.0065															19
6	.0049	.0076	.0105	.0134	.0164	.0195													12
8	.0081	.0127	.0175	.0224	.0274	.0325	.0377												24
10	.0120	.0188	.0259	.0332	.0407	.0483	.0560	.0638											16
12	.0166	.0261	.0359	.0459	.0562	.0667	.0774	.0882	.0991										24
14		.0343	.0471	.0604	.0739	.0877	.1017	.1159	.1303	.1449									21
16			.0597	.0765	.0937	.1112	.1289	.1470	.1652	.1836	.2022								16
18			.0736	.0943	.1155	.1370	.1589	.1811	.2036	.2263	.2493	.2724	.2958						18
20			.0888	.1137	.1392	.1652	.1916	.2184	.2455	.2729	.3005	.3284	.3566	.3850					11
22				.1347	.1649	.1957	.2270	.2586	.2907	.3232	.3559	.3890	.4223	.4559	.4898				10
24				.1572	.1925	.2284	.2649	.3019	.3393	.3771	.4154	.4540	.4929	.5321	.5716				13
26					.2219	.2633	.3053	.3479	.3911	.4347	.4788	.5233	.5681	.6134	.6589				6
28					.2531	.3003	.3482	.3969	.4461	.4959	.5461	.5969	.6480	.6996	.7515	.8039			10
30					.2860	.3394	.3936	.4486	.5042	.5605	.6173	.6746	.7325	.7908	.8495	.9086			7
32						.3806	.4414	.5030	.5654	.6285	.6922	.7565	.8214	.8868	.9526	1.019	1.086		10
34							.4916	.5602	.6297	.6999	.7709	.8425	.9147	.9875	1.061	1.135	1.209		8
36									.6969	.7747	.8532	.9325	1.012	1.093	1.174	1.256	1.338		3
38									.7671	.8527	.9392	1.026	1.114	1.203	1.292	1.382	1.473	1.564	4
40									.8403	.9340	1.029	1.124	1.221	1.318	1.416	1.514	1.613	1.713	4
42									.9163	1.019	1.122	1.226	1.331	1.437	1.544	1.651	1.759	1.868	3
44										1.106	1.218	1.332	1.446	1.561	1.677	1.793	1.911	2.029	2
46										1.197	1.318	1.441	1.564	1.689	1.814	1.941	2.068	2.196	2
48											1.422	1.554	1.687	1.821	1.957	2.093	2.230	2.368	0
50												1.671	1.814	1.958	2.104	2.250	2.398	2.546	0
52													1.945	2.100	2.255	2.412	2.570	2.729	1
54														2.245	2.412	2.580	2.749	2.919	1
56															2.573	2.752	2.932	3.113	1
58																	3.120	3.313	0
																			246

Note: Shading indicates range of data.

Data collected by destructive sampling of 246 trees in accessible locations of the Yukon.
 Table shows total volume of the stem, inside bark. The DBH shown is outside bark (field measurement).
 The height shown is total tree height. There is no allowance for defect.
 Table volumes were obtained by means of a logarithmic equation derived by the least squares method.
 Equation: Volume = .0000433 D^{1.77531} H^{1.10980}
 Standard error of estimate = .1420.
 Coefficient of determination (R²) = .994.

Table 2. Cubic metre volume table for white spruce (all ages), Yukon Territory, Ecoregions 4, 5, and 8.

DBH (cm)	Height (m)														Basis (No. of Trees)	
	4	6	8	10	12	14	16	18	20	22	24	26	28	30		
4	.0023	.0036	.0050	.0064												8
6	.0048	.0075	.0103	.0132	.0162											11
8	.0079	.0125	.0172	.0221	.0270	.0321										9
10	.0118	.0185	.0256	.0328	.0402	.0478	.0554									6
12	.0163	.0256	.0353	.0453	.0556	.0660	.0766	.0874								11
14	.0214	.0337	.0465	.0596	.0731	.0868	.1008	.1150								14
16		.0427	.0589	.0756	.0926	.1101	.1278	.1457	.1639							14
18		.0526	.0726	.0931	.1142	.1357	.1575	.1796	.2021							12
20			.0875	.1123	.1377	.1636	.1899	.2166	.2437	.2711						5
22				.1330	.1631	.1938	.2249	.2566	.2886	.3211						4
24				.1553	.1904	.2261	.2625	.2995	.3369	.3747	.4130					2
26					.2194	.2607	.3027	.3452	.3884	.4320	.4761	.5207				5
28					.2503	.2974	.3452	.3938	.4430	.4928	.5431	.5939				4
30					.2830	.3362	.3902	.4451	.5008	.5570	.6139	.6714				1
32					.3173	.3770	.4376	.4992	.5616	.6247	.6885	.7529				6
34						.4198	.4874	.5560	.6254	.6957	.7668	.8385				0
36							.5395	.6154	.6923	.7701	.8487	.9281	1.008			2
38							.5939	.6774	.7621	.8477	.9343	1.022	1.110			0
40								.7420	.8347	.9286	1.023	1.119	1.216	1.313		2
42									.9103	1.013	1.116	1.220	1.326	1.432		0
																116

Note: Shading indicates range of data.

Data collected by destructive sampling of 116 trees in accessible locations of the Yukon.

Table shows total volume of entire stem, inside bark. The DBH shown is outside bark (field measurement).

The height shown is total tree height. There is no allowance for defect.

Table volumes obtained by means of logarithmic equation derived by the least squares method.

$$\text{Equation: Volume} = .0000419 D^{1.77630} H^{1.11743}$$

Standard error of estimate = .0983.

Coefficient of determination (R^2) = .996.

Table 3. Cubic metre volume table for black spruce (all ages), Yukon Territory, Ecoregions 2, 5, 11, and 12.

DBH (cm)	Height (m)											Basis (No. of Trees)	
	4	6	8	10	12	14	16	18	20	22	24		
4	.0024	.0037	.0050										16
6	.0050	.0077	.0105	.0133									16
8	.0084	.0130	.0177	.0225	.0273	.0322							16
10	.0126	.0195	.0266	.0337	.0410	.0484	.0559						19
12	.0176	.0272	.0370	.0470	.0572	.0675	.0779	.0884					19
14		.0360	.0490	.0623	.0758	.0894	.1032	.1171	.1311				24
16		.0459	.0626	.0795	.0967	.1140	.1316	.1493	.1672	.1852			17
18		.0569	.0775	.0985	.1198	.1413	.1631	.1851	.2073	.2296	.2520		19
20		.0690	.0940	.1194	.1452	.1713	.1977	.2243	.2511	.2782	.3054		8
22			.1118	.1420	.1727	.2038	.2352	.2669	.2988	.3310	.3634		9
24			.1310	.1664	.2024	.2388	.2756	.3127	.3502	.3879	.4258		5
26				.1926	.2342	.2763	.3189	.3619	.4052	.4488	.4927		4
28					.2681	.3163	.3650	.4142	.4638	.5137	.5640		1
30							.4139	.4697	.5259	.5826			0
32								.5283	.5916				0
													173

Note: Shading indicates range of data.

Data collected by destructive sampling of 173 trees in accessible locations of the Yukon.

Table shows total volume of the stem, inside bark. The DBH shown is outside bark (field measurement).

The height shown is total tree height. There is no allowance for defect.

Table volumes obtained by means of logarithmic equation derived by the least squares method.

Equation: $\text{Volume} = .0000429 D^{1.82290} H^{1.07309}$

Standard error of estimate = .0972.

Coefficient of determination (R^2) = .996.

Table 4. Cubic metre volume table for black spruce (all ages), Yukon Territory, Ecoregions 4, 8, and 9.

DBH (cm)	Height (m)											Basis (No. of Trees)
	4	6	8	10	12	14	16	18	20	22	24	
4	.0024	.0036	.0048	.0060								8
6	.0052	.0078	.0103	.0129								7
8	.0089	.0133	.0177	.0221	.0265							13
10	.0135	.0202	.0270	.0337	.0404	.0470	.0537					13
12	.0190	.0285	.0380	.0474	.0568	.0662	.0757	.0851				25
14	.0254	.0381	.0507	.0633	.0759	.0885	.1010	.1136	.1262			23
16	.0327	.0489	.0651	.0813	.0975	.1137	.1298	.1460	.1621			10
18		.0610	.0812	.1014	.1216	.1418	.1619	.1821	.2022			8
20		.0743	.0990	.1236	.1482	.1728	.1973	.2219	.2464			7
22			.1184	.1478	.1772	.2066	.2360	.2653	.2947	.3240		3
24			.1394	.1740	.2087	.2433	.2778	.3124	.3469	.3815		6
26			.1620	.2023	.2425	.2827	.3229	.3630	.4032	.4433		2
28				.2324	.2787	.3249	.3711	.4172	.4633	.5094		2
30						.3698	.4224	.4749	.5274			0
32								.5360				0
												127

Note: Shading indicates range of data.

Data collected by destructive sampling of 127 trees in accessible locations of the Yukon.

Table shows total volume of entire stem, inside bark. The DBH shown is outside bark (field measurement).

The height shown is total tree height. There is no allowance for defect.

Table volumes obtained by means of logarithmic equation derived by the least squares method.

Equation: $\text{Volume} = .0000452 D^{1.87679} H^{0.99520}$

Standard error of estimate = .1059.

Coefficient of determination (R^2) = .993.

Table 5. Cubic metre volume table for lodgepole pine (all ages), Yukon Territory, Ecoregions 2 and 5.

DBH (cm)	Height (m)												Basis (No. of Trees)	
	4	6	8	10	12	14	16	18	20	22	24	26		
4	.0028	.0040	.0052	.0064	.0076									9
6	.0061	.0088	.0115	.0141	.0167	.0193								8
8	.0107	.0155	.0202	.0248	.0293	.0338	.0382							13
10	.0165	.0240	.0312	.0383	.0453	.0522	.0590	.0658						11
12	.0236	.0342	.0445	.0547	.0647	.0745	.0842	.0939	.1034					14
14		.0462	.0602	.0739	.0874	.1007	.1138	.1268	.1397	.1525				5
16			.0781	.0959	.1134	.1306	.1477	.1646	.1813	.1979	.2144			14
18				.1207	.1427	.1644	.1859	.2072	.2282	.2491	.2699	.2905		10
20				.1482	.1753	.2020	.2284	.2545	.2803	.3060	.3315	.3568		4
22				.1786	.2111	.2433	.2751	.3065	.3377	.3686	.3993	.4298		2
24				.2116	.2502	.2883	.3260	.3633	.4002	.4369	.4732	.5094		1
26				.2474	.2926	.3371	.3811	.4247	.4679	.5108	.5533	.5955		1
28				.2859	.3381	.3896	.4405	.4908	.5408	.5903	.6394	.6882		2
30					.3869	.4458	.5040	.5616	.6187	.6754	.7316	.7875		1
32						.5056	.5717	.6370	.7018	.7661	.8299	.8932		1
34							.6435	.7171	.7900	.8623	.9341	1.005		1
36								.8017	.8833	.9641	1.044			0
38										1.071				0
														97

Note: Shading indicates range of data.

Data collected by destructive sampling of 97 trees in accessible locations of the Yukon.
 Table shows total volume of the stem, inside bark. The DBH shown is outside bark (field measurement).
 The height shown is total tree height. There is no allowance for defect.
 Table volumes obtained by means of logarithmic equation derived by the least squares method.
 Equation: $\text{Volume} = .0000515 D^{1.95239} H^{0.91923}$
 Standard error of estimate = .1232.
 Coefficient of determination (R^2) = .993.

Table 6. Cubic metre volume table for lodgepole pine (all ages), Yukon Territory, Ecoregions 4 and 12.

DBH (cm)	Height (m)											Basis (No. of Trees)			
	4	6	8	10	12	14	16	18	20	22	24		26		
4	.0026	.0038	.0049												5
6	.0060	.0086	.0111	.0136	.0161	.0185									7
8	.0106	.0153	.0198	.0242	.0286	.0329	.0371								4
10	.0166	.0239	.0310	.0379	.0447	.0514	.0580	.0645							4
12	.0239	.0344	.0446	.0546	.0644	.0740	.0835	.0929							4
14	.0325	.0469	.0608	.0744	.0877	.1008	.1137	.1265	.1391						7
16		.0612	.0794	.0971	.1146	.1317	.1486	.1653	.1818						6
18			.1005	.1230	.1450	.1667	.1881	.2092	.2301						4
20				.1519	.1791	.2059	.2323	.2584	.2842						2
22				.1838	.2167	.2491	.2811	.3127	.3439	.3749					2
24				.2188	.2580	.2966	.3346	.3722	.4094	.4462					0
26				.2568	.3028	.3481	.3928	.4369	.4806	.5238					2
28				.2979	.3513	.4038	.4556	.5068	.5574	.6076					3
30				.3420	.4033	.4636	.5231	.5819	.6400	.6976	.7547				1
32				.3892	.4590	.5276	.5953	.6622	.7283	.7939	.8588				1
34					.5182	.5957	.6721	.7476	.8223	.8963					0
36						.6679	.7536	.8383	.9220						0
38															0
															52

Note: Shading indicates range of data.

Data collected by destructive sampling of 52 trees in accessible locations of the Yukon.

Table shows total volume of entire stem, inside bark. The DBH shown is outside bark (field measurement).

The height shown is total tree height. There is no allowance for defect.

Table volumes obtained by means of logarithmic equation derived by the least squares method.

$$\text{Equation: Volume} = .0000470 D^{2.00235} H^{0.90398}$$

Standard error of estimate = .0875.

Coefficient of determination (R^2) = .998.

Table 7. Cubic metre volume table for trembling aspen (all ages), Yukon Territory, Ecoregions 2, 4, 5, 9, 11, and 12.

DBH (cm)	Height (m)											Basis (No. of Trees)			
	4	6	8	10	12	14	16	18	20	22	24		26		
4	.0023	.0035	.0048	.0060	.0073										18
6	.0050	.0076	.0102	.0129	.0157	.0184									16
8	.0085	.0131	.0176	.0223	.0270	.0317	.0365								18
10	.0130	.0199	.0269	.0340	.0412	.0484	.0556	.0630							19
12		.0281	.0380	.0480	.0581	.0683	.0785	.0889							16
14		.0376	.0508	.0642	.0777	.0914	.1051	.1189	.1328						18
16		.0484	.0654	.0826	.1001	.1176	.1353	.1531	.1709						19
18		.0604	.0817	.1032	.1250	.1469	.1690	.1912	.2135	.2360					13
20		.0738	.0997	.1260	.1525	.1793	.2062	.2333	.2606	.2880					11
22			.1194	.1509	.1826	.2147	.2469	.2794	.3120	.3448	.3777				5
24			.1407	.1778	.2153	.2531	.2911	.3293	.3678	.4064	.4453				5
26				.2069	.2504	.2944	.3386	.3831	.4279	.4728	.5180				4
28				.2380	.2881	.3386	.3895	.4407	.4922	.5439	.5959				4
30				.2711	.3282	.3858	.4438	.5021	.5607	.6197	.6788				4
32					.3708	.4358	.5013	.5672	.6335	.7000	.7669				2
34					.4158	.4887	.5622	.6361	.7104	.7850	.8600				2
36					.4632	.5445	.6263	.7086	.7914	.8746	.9581				1
38						.6031	.6937	.7849	.8765	.9687					0
40							.7643	.8648	.9658						0
															175

Note: Shading indicates range of data.

Data collected by destructive sampling of 175 trees in accessible locations of the Yukon.

Table shows total volume of entire stem, inside bark. The DBH shown is outside bark (field measurement).

The height shown is total tree height. There is no allowance for defect.

Table volumes obtained by means of logarithmic equation derived by the least squares method.

Equation: $\text{Volume} = .0000392 D^{1.88989} H^{1.04839}$


Standard error of estimate = .0943.

Coefficient of determination (R^2) = .996.

APPENDIX A

Ecoregion	Number of Samples			
	White Spruce	Black Spruce	Lodgepole Pine	Trembling Aspen
1	68	-----	-----	-----
2	48	42	54	34
4	40	34	29	28
5	35	32	43	30
8	41	40	-----	-----
9	30	53	-----	20
11	47	47	-----	23
12	53	52	23	40
Total	362	300	149	175

APPENDIX B



From: Oswald and Senyk (1977)

APPENDIX C

Covariance Analysis: White Spruce

Source	Analysis of Covariance Sum of Squares	Degrees of Freedom (df)
Deviations from common line	6.15689	358
Deviations from group lines	4.59283	355
Due to differences between group lines	1.56406	3
	<u>Residual Sum of Squares</u>	
Ecoregion—(1, 2, 9, 11, 12)	3.52802	242
—(4, 5, 8)	1.06481	113
All regions	6.15689	358

H_A : Common Line

$$F = \frac{1.56406/3}{4.59283/355} = \frac{0.5214}{0.0129} = 40.4186 \quad (3, 355 \text{ df})$$

F significant at 1% level, i.e., group lines better than common line

Appendix C (continued)

Covariance Analysis: Black Spruce

Source	Analysis of Covariance Sum of Squares	Degrees of Freedom (df)
Deviations from common line	2.33066	297
Deviations from group lines	2.00660	294
Due to differences between group lines	0.32406	3
	<u>Residual Sum of Squares</u>	
Ecoregion—(2, 5, 11, 12)	1.34179	170
—(4, 8, 9)	0.66481	124
All regions	2.33066	297

H_A : Common Line

$$F = \frac{0.3241/3}{2.0066/294} = \frac{0.1080}{0.0068} = 15.8824 \text{ (3, 294 df)}$$

F significant at 1% level, i.e., group lines better than common line

Appendix C (continued)

Covariance Analysis: Lodgepole Pine

Source	Analysis of Covariance Sum of Squares	Degrees of Freedom (df)
Deviations from common line	1.96584	144
Deviations from group lines	1.80652	141
Due to differences between group lines	0.15932	3
	Residual Sum of Squares	
Ecoregion—(2, 5)	1.43350	93
—(4, 12)	0.37302	48
All regions	1.96584	144

H_A : Common Line

$$F = \frac{0.15932/3}{1.80652/141} = \frac{0.0531}{0.0128} = 4.1484 \text{ (3, 141 df)}$$

F significant at 1% level, i.e., group lines better than common line